

# Paltelet Count To Spleen Diameter Ratio As a Predictor of Esophageal Varices In Choronic Liver Diseases Due To Hepatitis C

Dr Muhammad Usman<sup>1</sup>, Dr Zeeshan Zafar<sup>2</sup>, Dr Zara Khatoon<sup>3</sup>, Dr Muhammad Asad Iqbal<sup>4</sup>, Dr Hafsa Ansar<sup>5</sup>, Dr Muhammad Nabeel Khan<sup>6</sup>, Dr Bilal Khursheed<sup>7</sup>

<sup>1</sup>DHQ Hospital Mirpur, Email: usmanchaudhary129@gmail.com
<sup>2</sup>DHQ Hospital Mirpur, Email: zeeshanstar102@gmail.com
<sup>3</sup>DHQ Hospital Mirpur, Email: jabeenzara.1996@gmail.com
<sup>4</sup>DHQ Hospital Mirpur, Email: drasadpharma418@gmail.com
<sup>5</sup>DHQ Hospital Mirpur, Email: hafsaansar.3913@gmail.com
<sup>6</sup>DHQ Hospital Mirpur, Email: nabeelkhan3621@gmail.com
<sup>7</sup>DHQ Hospital Mirpur, Email: drbilalkhursheed@gmail.com

ARTICLE INFO	ABSTRACT
Keywords: Platelet Count to Spleen Diameter Ratio, Esophageal Varices, Chronic Liver Disease, Hepatitis C. Corresponding Author:Dr Muhammad Usman, DHQ Hospital Mirpur, Email: usmanchaudhary129@gmail.com	<b>Objective:</b> To determine the diagnostic accuracy of platelet, count to spleen diameter ratio for presence of oesophageal varices in patients of chronic liver disease secondary to Hepatitis C taking endoscopy as gold standard. <b>Material and Methods:</b> The design of this study was cross sectional study. This study was carried out in Department of Medicine DHQ Hospital Mirpur and the duration of this study were 6 months from January 2024 to June 2024. The sampling technique of this study was non-probability consecutive sampling. After informed consent from patients, 180 patients fulfilling clinical case definition of CLD secondary to Hepatitis C were included in the study. Upper GIT endoscopy was done by standard protocol and oesophageal varices having a size of more than or equal to 5 millimetre (mm) were considered positive. The platelet counts in number per micro litre ( $\mu$ I) was measured by doing abdominal ultrasonography (USG). <b>Results:</b> The study population consisted of 180 patients of CLD secondary to HCV. Mean age of patients was 50.76±12.872 years. There were 86 (47.8%) male and 94 (52.2%) females, out of 180 patients. On Upper GIT endoscopy, the EVs were present in 95 (52.8%) and 85 (47.2%) patients have no EVs. The mean of PC/SD ratio of patients with varices was 31.0287 and those of without varices was 1358.4154. The mean of PC/SD ratio of all the patients was 1080.0724. The sensitivity and specificity for the diagnostic accuracy of PC/SD ratio were 71.6% and 64.9% respectively. The positive predictive value and negative predictive value were 66.0% and 64.9% respectively. The diagnostic accuracy of PC/SD ratio for prediction of EVs was 65.6%.

### **INTRODUCTION**

Chronic Liver Disease is a progressive and relapsing liver disease that occurs due to persistent injury to liver parenchyma by viral infection, toxins, deposition of copper in Wilson disease & iron in Hemochromatosis and autoimmune hepatitis. Amongst all the above causes of cirrhosis, Hepatitis C virus infection is common in Pakistan (1) Chronic Liver Disease (CLD) remains asymptomatic till the end stage liver disease occurs which manifests with complications of ascites, gastro esophageal variceal bleeding, encephalopathy, coagulopathy, spontaneous bacterial peritonitis and hepato-renal syndrome. WHO estimated that ESLD is responsible for the 1.1% of all deaths (2,3). Cirrhosis of the liver causes increase pressure to flow of blood in portal system. When portal pressure rises above 20 mmHg (3) it causes dilatation of veins at Portosystemic junctions which are situated at gastro-esophageal junctions, peri-umbilical area and ano-rectal junction. The dilated veins at gastro-esophageal junction are called gastro-esophageal varices. The gastro-esophageal varices are prevalent in 24-80% of ESLD patients (2). These varices bleed in 20-40% of patients (4) and mortality rate of variceal bleeding is 20-30% (3). According to the Baveno IV consensus 2005, every patient of cirrhotic liver should have screening endoscopy for early detection of esophageal varices (4). The patients having high risk esophageal varices (HREVs) should be given primary prophylaxis (non-selective  $\beta$  blockers or endoscopic band ligation) for the prevention of variceal bleeding which reduces risk of variceal bleeding up to 50% (4) But the American Association for the Study of Liver Disease (AASLD) recommends annual screening of varices by endoscopy in decompensated CLD and after every 2-3 years in compensated CLD, if varices are not present at index endoscopy. (5,6) To reduce burden on endoscopy unit for screening of varices, various studies are done to predict varices presence by non-invasive parameters (7). These variables include clinical radiological and biochemical markers, like liver enzymes serum albumin, platelet count, spleen diameter, portal vein diameter, liver span and presence of ascites (8). Although several studies are done to determine the diagnostic accuracy of PC/SD ratio for presence of esophageal varices there is variability among their results. Studies by Amin. K et al, Mattos AZ et al. and Sarwar (9). S et al have cut off value (909) of PC/SD ratio. Amin K et al showed significant diagnostic accuracy of PC/SD ratio for esophageal varices while Mattos AZ et al and Sara. S et al showed non-significant diagnostic accuracy of PC/SD ratio for esophageal varices (10), so further research is required to see diagnostic accuracy of PC/SD ratio for esophageal varices and to apply noninvasive parameters (PC/SD) for screening of esophageal varices.

### METHODOLOGY

The design of this study was cross sectional study. This study was carried out in Department of Medicine DHQ Hospital Mirpur and the duration of this study were 6 months from January 2024 to June 2024. 180 consecutive patients of Chronic liver disease secondary to Hepatitis C came in central endoscopy unit of Department of Medicine DHQ Hospital Mirpur via Emergency/OPD and had given informed consent for study were recruited for study. Detailed History was taken and thorough clinical examination was done of each patient. Platelet count (number/ micro liter) was done of each patient and spleen diameter was measured in millimeters by doing ultrasonography. The Upper gastro intestinal endoscopy of each patient was done by the experienced endoscopist for presence /absence of esophageal varices. All the information was collected on pre-designed proforma. The data is analyzed using SPSS statics.

#### RESULTS

The study population consisted of 180 cases of CLD secondary to HCV infection. The average age was  $50.82\pm12.912$  (mean ± Standard deviation). In the distribution of patients by gender, 86 (47.8%) patients were male and 94 (52.2%) patients were female. The means ± standard deviations of Platelet count (128116.67 ± 82727.747), spleen diameter (127.49 ± 24.758) and PC/SD ratio (1080.0724 ± 809.12154) are as shown in Table 1. There were 103 (57.2%) cases having PC/SD ratio less than 909, while 77(47.8%) have PC/SD ratio more than 909. The mean of PC/SD value having PC/SD ratio < 909 is 561.1746 ± 174.99 and those of PC/SD value > 909 is 1774.18 ± 804.86. Gastroscopy findings reveal 95(52.8%) cases have EVs present and there are no EVs in 85(47.2%) cases. The mean of spleen diameter in patients having EVs is 131.56 ± 24.285 and those, without EVs is 122.94 ± 24.627. The mean value of PC/SD ratio in patients having EVs on Gastroscopy is 831.0287 ± 594.32062 and in patients without EVs it is 1358.4154 ± 922.64748.

By analyzing the data of EVs on basis of PC/SD, taking Gastroscopy as gold standard test for detection of EVs, into 2 X 2 Table, the following results were observed. There were 68 true positive (TP) cases, 35 false positive (FP) cases, 50 true negative (TN) and 27 false negative (FN) cases. The sensitivity and specificity of PC/SD ratio as a predictor of esophageal are 71.6% and 58.8% respectively (Table 02). The positive predictive value and negative predictive value of this test is 66% and 64.9% respectively (Table 03) The positive likelihood ratio (LR+) of this test is 1.74 and negative likelihood ratio is 0.48. The diagnostic accuracy of PC/SD ratio is 65.6%. Summary of statistical analysis has been given in Table 04.

Sr. No	Variables	Range	Minimum	Maximum	Mean	Std. Deviation
1	Patient Age in Years	67	18	85	50.82	12.912
2	Platelet Count Per Micro Liter	411000	23000	434000	128116.67	82727.747
3	Spleen Diameter in Millimeter	130	80	210	127.49	24.758
4	Pc/Sd Ratio Value	4175.72	164.28	4340.00	1080.0724	809.12154

# Table 1- Descriptive Statistics of Numeric Data

#### Table .02- Esophageal Varices on Platelet Count to Spleen Diameter Ratio X Esophageal Varices on Gastroscopy Cross Tabulation Showing Sensitivity and Specificity

		Esophagea Gastre			
			> Present on Gastroscopy	Absent on Gastroscopy	Total
	PCSD less than		68	35	103
Esophageal varices on platelet count to spleen diameter ratio	909	% within esophageal varices on platelet count to spleen diameter ratio	71.6%	41.2%	57.2%
	PCSD more than		27	50	77
	909	% within esophageal varices on platelet count to spleen diameter ratio	28.4%	58.8%	42.8%
Total			95	85	180
		% within esophageal	100.0%	100.0%	100.0%
		varices on platelet count to spleen diameter ratio			

 Table 03-Esophageal Varices on Platelet Count to Spleen Diameter Ratio X Esophageal Varices on Gastroscopy

 Cross Tabulation Showing Positive Predictive Value and Negative Predictive Value

			Fsonhageal	varices on	Total
			Gastro	scopy	
			Gastroscopy	Gastroscopy	
	PCSD less than	Count	68	35	103

		s	ices on platelet count to spleen diameter ratio			
			Count	27	50	77
PCSD	more than 909	% within esophageal varices on platelet count to spleen diameter ratio		35.1%	64.9%	100.0%
Total				95	85	180
		<b>N</b> 1	% within esophageal varices on platelet count to spleen diameter ratio	52.8%	47.2%	100.0%

#### TABLE 04: Summary Of Statistical Analysis

Data	Value
Sensitivity	71.6%
Specificity	58.8%
Positive Predictive Value	66%
Negative Predictive Value	64.9%
Diagnostic Accuracy	65.6%.
Positive Likelihood Ratio (LR+)	1.74
Negative Likelihood Ratio (LR-)	0.48

#### DISCUSSION

Upper GIT bleeding is a renowned complication of CLD. In Pakistan, HCV infection is a most common cause of CLD (11). To reduce the risk of mortality caused by upper GIT bleeding, Baveno IV consensus on endoscopy updates recommended that every person of CLD should have screening by Gastroscopy at the time of diagnosis for the early detection of EVs and institution of primary prophylaxis once the EVs are present on Gastroscopy (12). As the Gastroscopy is an invasive procedure, having a risk of infection spread and also cost effectiveness and availability problems, noninvasive parameters like (platelets count and spleen diameter ratio) are studied by different researchers (2-7) to see the diagnostic accuracy for prediction of EVs including this study. The PC/SD ratio uses two parameters, which are routinely done in cirrhotic patients. In this study the mean age, of patients included, is 50.82 which is comparable to both of the studies by Amin. K et al (57.61) (3) and Mattos et al (56.6) (6). A female to male ratio is 1.09:1 with an overall female preponderance of 52% which is consistent with the study by Amin. K et al where female to male ratio was 1.03:1 (13).

In the present study, the EVs diagnosed by Gastroscopy are 52.8%, which fall in the range of 24-80% as mentioned in the literature. This reveals that about half of the patients, whose screening endoscopy is done, do not require endoscopy for early detection of EVs. In other words, they are exposed to infections by doing an invasive procedure. Noninvasive parameters can be established for detection of EVs to prevent the risks and discomfort of Gastroscopy (14,15). PC/SD ratio has gained more attention by the researchers as compared to other noninvasive parameters because of the fact that with time platelets count decreases in natural course of CLD, mainly due to hyper-splenism, immune-mediated destruction of platelets (in viral hepatitis) and decreased production of platelets secondary to myelo-toxic effect of viral hepatitis & alcohol and decreased production of thrombo-poietin (16). The spleen diameter increases during natural course of CLD secondary to portal hypertension. This study is based on the concept that PC/SD decreases during natural course of disease (due to decrease in platelet count and increased spleen size) along with complications of CLD, like EVs. Thus in patients with

decrease in PC/SD ratio the risk of development of EVs increases. A cut off value of 909 is used in this study, as used by Amin. K et al (3), Mattos et al (6) and Sarwar. Set (7) al in the literature. In this study, 68 (71.6%) patients having PC/SD ratio < 909 had EVs on Gastroscopy and 35 (41.1%) patients do not have EVs on Gastroscopy. There were 50(58.8%) patients having PC/SD Ratio >909 and no EVs on Gastroscopy, while 27 (28.8%) with PC/SD > 909 have EVs on Gastroscopy. By using the confidence interval (CI) of 95%, PC/SD ratio has sensitivity of 71.6%, specificity of 58.8%, PPV of 66%, NPV of 64.4% and diagnostic accuracy of 65.6%. The diagnostic accuracy of PC/SD is comparable to that predicted by Mattos et al (68.9%) (6). Although this study reveals association of PC/SD ratio with presence of EVs on Gastroscopy, but its sensitivity, specificity, PPV, NPV and diagnostic accuracy are not same as, are described by Amin. K et al (3). Our study does not favor the use of PC/SD ratio for prediction of EVs because the risk of missing the patients having EVs is more dangerous, leading to the risk of bleeding from EVs which has greater mortality in patients of CLD (4). Moreover, a cut off value of (909) for detection of EVs should be re-evaluated as advised by Mattos. AZ et al (6) Sarwar. S et al (7) and Schwarzenegger. E et al (8).

## CONCLUSION

We are not in favor of platelet count/spleen diameter ratio to be used as an adequate index for the prediction of EVs in cirrhotic patients. At the moment, we still think cirrhotic patients must have a screening endoscopy in order to identify EVs at the time of their diagnosis. It is also important that PC/SD ratio having a different cut off value and other non-invasive parameters should be developed in order to reduce the burden on health system and to improve cost effectiveness for early detection of EVs and initiation of primary prophylaxis for EVs.

#### REFERENCES

- 1. Jafri. W, Subhan. A. Hepatitis C in Pakistan: Magnitude, Genotype, Disease Characteristics and Therapeutic Response. Tropical Gastroenterol 2022; 29(4):194–201.
- 2. Abu El Makarem MA, Shatat ME, Shaker Y, Abdel Aleem AA, El Sherif AM, Abdel Moaty M, et al. Platelet count/bipolar spleen diameter ratio for the prediction of esophageal varices: The special Egyptian situation. Hepat Mon 2021; 11(4):278-84.
- 3. Amin k. et al.Platelet Count to Spleen diameter ratio as apredictor of esophageal varices in patients of liver cirrhosis due to Hepatitis C virus Infection. JUMDC 2022; 03(1):06-11.
- 4. Barrera F. et al. Platelet count/bipolar spleen diameter ratio for non-invasive prediction of high risk esophageal varices in cirrhotic patients. Annals of hepatol 2021; 08(04):325-330.
- 5. Krystallis C et al. Update of endoscopy in liver disease: More than just treating varices. World J Gastroenterol 2022; 18(5): 401-411.
- 6. Mattos AZ, Mattos AA, Vianna FF, Musskopf MI, Pereira-Lima JC, Maciel AC. Platelet count/spleen diameter ratio: analysis of its capacity as a predictor of the existence of esophageal varices. Arq Gastroenterol 2020; 47(3):275-78.
- 7. Sarwar S, Alam A, Khan AA, Butt AK, Shafqat F, Shah WH et al. Platelet count/splenic size ratio: Can it predict the presence of varices in patients of Cirrhosis of liver? Proceeding Sheikh Zayed Postgrad Med Inst 2024; 18: 21-61.
- 8. Schwarzenegger E, Meyer T, Golla V, Sahdala NP, Min AD. Utilization of platelet count spleen diameter ratio in predicting the presence of esophageal varices in patients with cirrhosis. J Clin Gastroenterol. 2022; 44(2):146-50.
- 9. Mossie, G. Y., Nur, A. M., Ayalew, Z. S., Azibte, G. T., & Berhane, K. A. (2024). Platelet counts to spleen diameter ratio: A promising noninvasive tool for predicting esophageal varices in cirrhosis patients. World Journal of Hepatology, 16(10), 1177.
- 10. Yu, S., Chen, W., & Jiang, Z. (2021). Platelet count/spleen volume ratio has a good predictive value for esophageal varices in patients with hepatitis B liver cirrhosis. PLoS One, 16(12), e0260774.
- 11. GR, S., & Kulkarni, S. (2022). " Comparative study of the accuracy of Platelet count/spleen diameter ratio with other non-invasive parameters in predicting large esophageal varices in patients with chronic liver disease". Medica Innovatica, 11(2).
- 12. ElDesoky, A. E., Aeriba, M., Abd ElFattah, F., ElHawary, G., & Awad, M. M. (2022). Platelet count/spleen diameter ratio as a predictor of high-risk esophageal varices in patients with liver cirrhosis. Medical Journal of Viral Hepatitis, 6(2), 25-30.
- 13. Patil, S., Patnaik, S. K., Kanungo, M., Uthansingh, K., Narayan, J., Pradhan, S., ... & Pati, G. K. (2024). Platelet Count/Spleen Diameter Ratio as a Non-Invasive Predictor of Esophageal Varices in Cirrhotic Patients: A Single-Center Experience. Gastroenterology Insights, 15(1), 98-106.

- 14. Ajay Mal Reventh, P. (2022). A Prospective study on Predicting Esophageal Varices in Cirrhosis with Platelet Count to Spleen Diameter Ratio, Right Lobe of Liver Diameter to Serum Albumin Ratio, Portal Vein Diameter in a Tertiary Care Hospital in Chennai (Doctoral dissertation, Madras Medical College, Chennai).
- 15. Ajay Mal Reventh, P. (2022). A Prospective study on Predicting Esophageal Varices in Cirrhosis with Platelet Count to Spleen Diameter Ratio, Right Lobe of Liver Diameter to Serum Albumin Ratio, Portal Vein Diameter in a Tertiary Care Hospital in Chennai (Doctoral dissertation, Madras Medical College, Chennai).
- 16. Sridhar, S. (2023). A Prospective Study on Predicting Esophageal Varices in Cirrhosis with Platelet Count to Spleen Diameter Ratio, Right Lobe of Liver Diameter to Serum Albumin Ratio, Portal Vein Diameter in A Tertiary Care Hospital in Chennai. International Journal, 6(3), 247.
- 17. Suman, G. R., & Sriharsha Kulkarni, I. G. (2022). Comparative study of the accuracy of Platelet count/spleen diameter ratio with other non-invasive parameters in predicting large esophageal varices in patients with chronic liver disease. Medica, 11(2), 69.